information from herbarium material I have become aware of several locations of this plant throughout New England and elsewhere. If anyone has any information regarding *Podostemum ceratophyllum*, or knows of any populations, I would be most pleased to hear of them.

LITERATURE CITED

- Accorsi, W. R. 1950. Biology and Ecology of the Podostemaceae of the Piracicaba Fall. Proc. Seventh Int. Bot. Congr. Stockholm, 1950. 681-682.
- HAMMOND, B. L. 1936. Regeneration of Podostemum ceratophyllum. Bot. Gaz. 97: 834-845.
- _____ 1937. Development of *Podostemum ceratophyllum*. Bull. Torrey Bot. Club **64**: 17-36.
- LAWRENCE, G. H. M. 1951. Taxonomy of Vascular Plants. The MacMillan Co., N.Y.
- ROYEN, P. VAN. 1951. The Podostemaceae of the New World. I. Meded. Bot. Mus. Utrecht 107: 1-151.
- Went, F. A. F. C. 1926. Morphological and Histological Peculiarities of the Podostemaceae. Proc. Intern. Congr. Pl. Sci. Ithaca. Vol. I: 351-358.
- WILLIS, J. C. 1902. Studies on the Morphology and Ecology of the Podostemaceae of Ceylon and India. Ann. Roy. Bot. Gardens, Peradeniya Vol. I(4): 267-465.

DEPARTMENT OF BOTANY AND PLANT PATHOLOGY NESMITH HALL UNIVERSITY OF NEW HAMPSHIRE DURHAM, N. H. 03824

THE REDISCOVERY OF JUNCUS STYGIUS IN MAINE

C. BARRE HELLQUIST

Juncus stygius L. var. americanus Buchenau is a northern bog rush found throughout Canada and rarely into northern United States from Maine to Minnesota. This rush is on the rare and endangered lists of Maine (Eastman, 1978), New York (Mitchell et al. 1980) and Michigan (Wagner et al., 1977). A total of five locations have been reported for this species from Minnesota (Wheeler & Glaser, 1979). Three historical records are known from Maine: two from Fort Kent and one from Crystal Bog, Crystal. The Fort Kent populations were last reported in 1908. The Crystal Bog population was discovered in

1907 by M. L. Fernald. Tyler and Gawler (1980) indicated that this rush had not been found recently.

During August 1979, while doing field work with Harry Tyler, Jr. and Kathy Kehoe on the wetland areas in Aroostook County, Garrett Crow of the University of New Hampshire and I were taken to the bog. Accompanying us were Candy McKellar and Sally Rooney, of the Nature Conservancy's Crystal Bog Stewardship Committee, who have been conducting botanical investigations of the bog. They stated at that time that *Juncus stygius* was the only plant not rediscovered in the bog. Upon entering the bog I spotted several plants and then found it to be common. The site was the large open portion of the bog west of the *Thuja* stand. Here *J. stygius* was found growing with *Rhynchospora alba* (L.) Vahl., with which it might be confused. This species can easily be overlooked and may occur more frequently in other bogs of Aroostook County. Specimens have been deposited in the herbaria of Boston State College, The University of Maine, and the New England Botanical Club.

LITERATURE CITED

- Eastman, L. M. 1978. Rare and endangered vascular plant species in Maine. The New England Botanical Club in cooperation with the U. S. Fish and Wildlife Service. 33 pp.
- MITCHELL, R. S., C. J. SHEVIAK, & J. K. DEAN. 1980. Rare and endangered vascular plant species in New York State. The State Botanist's Office, N.Y. State Museum, Albany in cooperation with the U. S. Fish and Wildlife Service. 38 pp.
- TYLER, H. R., JR., & S. G. GAWLER. 1980. The botanical aspect of Maine's Critical Areas Program. Rhodora 82: 207-225.
- WAGNER, W. H., E. G. Voss, J. H. BEAMAN, E. A. BOURDO, F. W. CASE, J. A. CHURCHILL, & P. W. THOMPSON. 1977. Endangered, threatened, and rare vascular plants in Michigan. Mich. Bot. 16: 99-122.
- WHEELER, G. A., & P. H. GLASER. 1979. Notable vascular plants of the Red Lake peatland, northern Minnesota. Mich. Bot. 18: 137-142.

DEPARTMENT OF BIOLOGY BOSTON STATE COLLEGE 625 HUNTINGTON AVENUE BOSTON, MA. 02115